

ON THE PRE-BÉZOUT PROPERTY OF WIENER ALGEBRAS ON THE DISC AND THE HALF-PLANE

RAYMOND MORTINI AND AMOL SASANE

ABSTRACT. Let \mathbb{D} denote the open unit disk $\{z \in \mathbb{C} \mid |z| < 1\}$, and \mathbb{C}_+ denote the right half-plane $\{s \in \mathbb{C} \mid \operatorname{Re}(s) \geq 0\}$.

- (1) Let $W^+(\mathbb{D})$ be the Wiener algebra of the disc, that is the set of all absolutely convergent Taylor series in the open unit disk \mathbb{D} , with pointwise operations.
- (2) Let $W^+(\mathbb{C}_+)$ be the set of all functions defined in the right half-plane \mathbb{C}_+ that differ from the Laplace transform of a function $f_a \in L^1(0, \infty)$ by a constant. Equipped with pointwise operations, $W^+(\mathbb{C}_+)$ forms a ring.

We show that the rings $W^+(\mathbb{D})$ and $W^+(\mathbb{C}_+)$ are pre-Bézout rings.

CDAM Research Report LSE-CDAM-2008-4

1. INTRODUCTION

The aim of this paper is to show that the rings $W^+(\mathbb{D})$ and $W^+(\mathbb{C}_+)$ (defined below) are pre-Bézout.

We first recall the notion of a pre-Bézout ring.

Definition 1.1. Let R be a commutative, unital ring.

- (1) An element $d \in R$ is called a *greatest common divisor* of $a, b \in R$ if it is a divisor of a and b and if k is another divisor, then k divides d .
- (2) The ring R is said to be *pre-Bézout* if for every $a, b \in R$ for which there exists a greatest common divisor d , there exist $x, y \in R$ such that $d = xa + yb$.

Michael von Renteln [12, Theorem 2.4, p. 54] proved that the disc algebra $A(\mathbb{D})$ (the ring of continuous functions on the closed unit disc $\overline{\mathbb{D}}$, which are analytic in the open unit disc \mathbb{D} , with the usual pointwise operations) is pre-Bézout. The first author of the present paper [7] showed the pre-Bézout property for the Sarason algebra $QA = (C(\mathbb{T}) + \widehat{C(\mathbb{T})}) \cap H^\infty(\mathbb{D})$ of bounded analytic functions having quasicontinuous boundary values. (Here $\widehat{C(\mathbb{T})}$ denotes the set of harmonic conjugates of continuous functions on \mathbb{T} .) Note that the algebra $H^\infty(\mathbb{D})$ (of all bounded and analytic functions in the open unit disc, with pointwise operations) is not pre-Bézout [12, Remark, p. 54].

1991 *Mathematics Subject Classification.* Primary 30H05 ; Secondary 42A99, 93D15.
Key words and phrases. pre-Bézout ring, Wiener algebra.

In this article, we will show that the rings $W^+(\mathbb{D})$ and $W^+(\mathbb{C}_+)$ (defined below) are pre-Bézout.

Throughout the article, we will use the following notation:

$$\begin{aligned}\mathbb{D} &:= \{z \in \mathbb{C} \mid |z| < 1\} \\ \overline{\mathbb{D}} &:= \{z \in \mathbb{C} \mid |z| \leq 1\} \\ \mathbb{C}_+ &:= \{s \in \mathbb{C} \mid \operatorname{Re}(s) \geq 0\}.\end{aligned}$$

Definition 1.2.

- (1) The *Wiener algebra of the disc*, $W^+(\mathbb{D})$, is the set of all functions $f : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ such that f is analytic in \mathbb{D} and $\sum_{n=0}^{\infty} |a_n| < \infty$ for $f(z) = \sum_{n=0}^{\infty} a_n z^n$ ($z \in \mathbb{D}$). Equipped with pointwise operations and the norm $\|f\|_{W^+} := \sum_{n=0}^{\infty} |a_n|$, $W^+(\mathbb{D})$ is a Banach algebra.
- (2) Let $W^+(\mathbb{C}_+)$ denote the set of all functions $F : \mathbb{C}_+ \rightarrow \mathbb{C}$ such that $F(s) = \widehat{f}_a(s) + f_0$ ($s \in \mathbb{C}_+$), where $f_a \in L^1(0, \infty)$, $f_0 \in \mathbb{C}$, and \widehat{f}_a denotes the Laplace transform of f_a given by

$$\widehat{f}_a(s) = \int_0^{\infty} e^{-st} f_a(t) dt, \quad s \in \mathbb{C}_+.$$

Equipped with pointwise operations and the norm

$$\|F\|_{W^+} = \|f_a\|_{L^1} + |f_0|,$$

$W^+(\mathbb{C}_+)$ is a Banach algebra.

We note that $W^+(\mathbb{C}_+)$ is contained in the set of all holomorphic functions on the right half-plane that admit continuous extensions to the imaginary axis and have a limit at infinity. We will call $W^+(\mathbb{C}_+)$ the *Wiener-Laplace algebra*.

Remark 1.3.

- (1) From the application point of view, the above algebras also arise as a natural classes of transfer functions of stable distributed parameter systems in control theory; see [10].
- (2) We use the notation $W^+(\mathbb{C}_+)$ in order to highlight the similarity with $W^+(\mathbb{D})$. Indeed, $W^+(\mathbb{D})$ is isomorphic to the algebra of summable sequences $\ell^1(\mathbb{N})$ with convolution, pointwise addition, and the $\ell^1(\mathbb{N})$ norm. Now instead of this “discrete” convolution algebra, we consider “distributed” summable functions $L^1(0, \infty)$, again with convolution, pointwise addition, and the $L^1(0, \infty)$ -norm, and attach the identity element δ (=Dirac distribution) to it, we obtain the convolution algebra $L^1(0, \infty) + \mathbb{C}\delta$. Then $W^+(\mathbb{C}_+)$ is isomorphic to the algebra $L^1(0, \infty) + \mathbb{C}\delta$ via Laplace transformation.

Our main results are the following:

Theorem 1.4. *The ring $W^+(\mathbb{D})$ is pre-Bézout.*

Theorem 1.5. *The ring $W^+(\mathbb{C}_+)$ is pre-Bézout.*

Remark 1.6.

- (1) The relevance of the pre-Bézout property in control theory is the following:

Suppose R is a pre-Bézout ring and we have a plant whose transfer function p belongs to the field of fractions of R . Then p has a weakly coprime factorization iff p has a coprime factorization; see [9, Proposition, p. 54].

- (2) We recall that a commutative ring R is called *Bézout* if every finitely generated ideal in R is principal.

Neither of our algebras $W^+(\mathbb{D})$ nor $W^+(\mathbb{C}_+)$ are Bézout. That $W^+(\mathbb{D})$ is not Bézout can be shown by considering the ideal (f, g) , where

$$f = (1 - z)^3 \quad \text{and} \quad g = (1 - z)^3 e^{-\frac{1+z}{1-z}};$$

see [8, Remark after Theorem 1, p. 224]. On the other hand the fact that $W^+(\mathbb{C}_+)$ is not Bézout follows from a general result which says that if R is any subring of the ring H^∞ (of bounded analytic functions in the open right half-plane $\operatorname{Re}(s) > 0$, with pointwise addition and multiplication), such that R contains the Laplace transforms of functions from $L^1(0, \infty)$, then R has a finitely generated ideal which is not principal; see [6, Theorem].

In Sections 3 and 4 we will give the proofs of Theorems 1.4 and 1.5, respectively, but before doing that, in Section 2, we first give a few preliminaries.

2. PRELIMINARIES

It is well known that the maximal ideals (or kernels of multiplicative linear functionals) of $W^+(\mathbb{D})$ have the form

$$\mathfrak{m}_a = \{f \in W^+(\mathbb{D}) \mid f(a) = 0\}$$

for some $a \in \overline{\mathbb{D}}$. Similarly, the set of maximal ideals in $W^+(\mathbb{C}_+)$ coincides with the set of ideals of the form \mathfrak{M}_{s_0} and \mathfrak{M}_∞ , where

$$\mathfrak{M}_{s_0} = \{F \in W^+(\mathbb{C}_+) \mid F(s_0) = 0\}, \quad s_0 \in \mathbb{C}_+,$$

and \mathfrak{M}_∞ is given by the kernel of the homomorphism $\varphi : W^+(\mathbb{C}_+) \rightarrow \mathbb{C}$ defined by

$$F = \widehat{f}_a + f_0 \mapsto f_0 \quad (f_a \in L^1(0, \infty), f_0 \in \mathbb{C}).$$

That is,

$$\mathfrak{M}_\infty = \{F \in W^+(\mathbb{C}_+) \mid \exists f_a \in L^1(0, \infty) \text{ such that } F = \widehat{f}_a\} = \widehat{L^1(0, \infty)}.$$

Since every maximal ideal is closed, all the sets \mathfrak{m}_α , $|\alpha| = 1$, are commutative Banach subalgebras of $W^+(\mathbb{D})$. Similarly, $\mathfrak{M}_{i\beta}$, $\beta \in \mathbb{R}$, and \mathfrak{M}_∞ are commutative Banach subalgebras of $W^+(\mathbb{C}_+)$. Obviously these algebras have no identity element. But there is a substitute, namely the notion of the bounded approximate identity, which will be useful in the sequel.

Definition 2.1. Let R be a commutative Banach algebra (without identity element). We say that R has a *bounded approximate identity* if there exists a bounded sequence $(e_n)_n$ of elements e_n in R such that for any $f \in R$,

$$\lim_n \|e_n f - f\| = 0.$$

We will also need the following technical result:

Proposition 2.2 (Varopoulos, [14]). *Let R be a Banach algebra with a bounded left approximate identity. Then for every sequence $(a_n)_{n \geq 1}$ in R converging to 0, there exists a sequence $(b_n)_{n \geq 1}$ in R converging to 0, as well as an element $c \in R$ such that for all $n \geq 1$, $a_n = cb_n$.*

Lemma 2.3. *Let R be a commutative integral domain with identity 1. If d ($\neq 0$) is a greatest common divisor of f_1, \dots, f_n , then 1 is a greatest common divisor of $\frac{f_1}{d}, \dots, \frac{f_n}{d}$.*

Proof. Clearly 1 divides $\frac{f_1}{d}, \dots, \frac{f_n}{d}$. If h is a divisor of $\frac{f_1}{d}, \dots, \frac{f_n}{d}$, then $\frac{f_k}{d} = hg_k$, for some $g_k \in R$, $k = 1, \dots, n$. So dh is common divisor of f_1, \dots, f_n , and as d is the greatest common divisor of f_1, \dots, f_n , dh divides d , that is, $dhk = d$ for some $k \in R$. Since R is an integral domain and $d \neq 0$, we obtain $hk = 1$, that is, h divides 1, proving the claim. \square

3. $W^+(\mathbb{D})$ IS PRE-BÉZOUT

Let $z_0 \in \mathbb{T} := \{z \in \mathbb{D} \mid |z| = 1\}$. Consider the maximal ideal

$$\mathfrak{m}_{z_0} := \{f \in W^+(\mathbb{D}) \mid f(z_0) = 0\}.$$

We will use the following result on the existence of a bounded approximate identity for \mathfrak{m}_{z_0} . Without loss of generality, we take $z_0 = 1$.

Proposition 3.1 (Faivyševskij, [2]). *Let $n \in \mathbb{N}$, $(r_n)_{n \in \mathbb{N}}$ be any sequence such that $r_n \searrow 1$, and*

$$e_n(z) := \frac{z - 1}{z - r_n}.$$

Then $(e_n)_{n \in \mathbb{N}}$ is a bounded approximate identity for \mathfrak{m}_1 .

A rather lengthy proof of the above result in the case when $r_n = 1 + \frac{1}{n}$ can be found in [5, Lemma 1]. For the reader's convenience we present a short proof here.

Proof. A simple calculation gives that $\|\frac{z-1}{z-1-\epsilon}\|_{W^+} \leq 2$. Since the partial sums $S_n - S_n(1)$ for f approximate $f \in \mathfrak{m}_1$, it suffices to consider $q(z) = (z - 1)p(z)$, where $p \in \mathbb{C}[z]$. But

$$\left\| \frac{z-1}{z-1-\epsilon} q - q \right\|_{W^+} = \left\| \epsilon \frac{q}{z-1-\epsilon} \right\|_{W^+} = \epsilon \left\| \frac{z-1}{z-1-\epsilon} p \right\|_{W^+} \leq 2\epsilon \|p\|_{W^+}.$$

\square

We will also use the following fact proved on page 301 of the Proof of the Theorem in [13].

Proposition 3.2 (M. von Renteln). *Let $f \in W^+(\mathbb{D})$ and $z_0 \in \mathbb{D}$ be such that $f(z_0) = 0$. Then $\frac{f}{z-z_0} \in W^+(\mathbb{D})$.*

We will also need the corona theorem for $W^+(\mathbb{D})$; see for example [13, Theorem]:

Proposition 3.3. *If $f_1, \dots, f_n \in W^+(\mathbb{D})$ are such that*

$$\text{for all } z \in \overline{\mathbb{D}}, \quad |f_1(z)| + \dots + |f_n(z)| > 0,$$

then there exist $g_1, \dots, g_n \in W^+(\mathbb{D})$ such that

$$\text{for all } z \in \overline{\mathbb{D}}, \quad g_1(z)f_1(z) + \dots + g_n(z)f_n(z) = 1.$$

Lemma 3.4. *Suppose that $f_1, \dots, f_n \in W^+(\mathbb{D})$ and d is a greatest common divisor of f_1, \dots, f_n . If $z_0 \in \overline{\mathbb{D}}$ is a common zero of f_1, \dots, f_n , then $d(z_0) = 0$ as well.*

Proof. If $z_0 \in \mathbb{D}$, then let m be the least integer among the multiplicities of z_0 as a zero respectively of f_1, \dots, f_n . By Proposition 3.2, $(z - z_0)^m$ is a divisor of f_1, \dots, f_n . But since d is the greatest common divisor of f_1, \dots, f_n , it follows that $(z - z_0)^m$ divides d .

If on the other hand $z_0 \in \mathbb{T}$, then $f_1, \dots, f_n \in \mathfrak{m}_{z_0}$, where

$$\mathfrak{m}_{z_0} := \{f \in W^+(\mathbb{D}) \mid f(z_0) = 0\}.$$

By Proposition 3.1, \mathfrak{m}_{z_0} has a bounded approximate identity. Applying Proposition 2.2, with $(a_n)_{n \geq 1} := (f_1, \dots, f_n, 0, 0, 0, \dots)$, we get the existence of an element $c \in \mathfrak{m}_{z_0}$, and $g_1, \dots, g_n \in \mathfrak{m}_{z_0}$ such that $f_k = cg_k$, $k = 1, \dots, n$. So we have a common divisor c of f_1, \dots, f_n . Since $c(z_0) = 0$, and d is a greatest common divisor, we have that c divides d and hence $d(z_0) = 0$, too. \square

Proof of Theorem 1.4. Let $f_1, \dots, f_n \in W^+(\mathbb{D})$ have a greatest common divisor d ($\neq 0$). By the algebraic result in Lemma 2.3, it follows that 1 is a greatest common divisor of $\frac{f_1}{d}, \dots, \frac{f_n}{d}$. Lemma 3.4 gives

$$\left| \frac{f_1}{d} \right| + \dots + \left| \frac{f_n}{d} \right| > 0 \quad \text{in } \overline{\mathbb{D}}.$$

By Proposition 3.3, it follows that there exist $g_1, \dots, g_n \in W^+(\mathbb{D})$ such that

$$g_1 \frac{f_1}{d} + \dots + g_n \frac{f_n}{d} = 1,$$

and so $g_1 f_1 + \dots + g_n f_n = d$, completing the proof of the theorem. \square

4. $W^+(\mathbb{C}_+)$ IS PRE-BÉZOUT

We will first prove that the maximal ideals $\mathfrak{M}_{i\beta}$, $\beta \in \mathbb{R}$, and \mathfrak{M}_∞ in $W^+(\mathbb{C}_+)$ have a bounded approximate identity. To this end, we need the following lemma.

Lemma 4.1. *Suppose $F \in \mathfrak{M}_0$. Then for each $\epsilon > 0$, there exists $P \in \mathfrak{M}_0$ such that $P = \widehat{p}_a + p_0$, where $p_a \in L^1(0, \infty)$ has compact support, $p_0 \in \mathbb{C}$, and $\|F - P\|_{W^+} < \epsilon$.*

Proof. Let $\epsilon > 0$ be given. Let $F = \widehat{f}_a + f_0$, where $f_a \in L^1(0, \infty)$ and $f_0 \in \mathbb{C}$. Choose a compactly supported $p_a \in L^1(0, \infty)$ such that

$$\|p_a - f_a\|_{L^1} < \frac{\epsilon}{2}.$$

Set

$$P := \widehat{p}_a + \underbrace{\left(-\int_0^\infty p_a(t)dt\right)}_{=:p_0}.$$

Then $P \in W^+(\mathbb{C}_+)$ and

$$P(0) = \widehat{p}_a(0) + p_0 = \int_0^\infty p_a(t)dt - \int_0^\infty p_a(t)dt = 0.$$

So $P \in \mathfrak{M}_0$. We have

$$\begin{aligned} \left|f_0 + \int_0^\infty p_a(t)dt\right| &= \left|f_0 + \int_0^\infty f_a(t)dt + \int_0^\infty (p_a(t) - f_a(t))dt\right| \\ &= \left|f_0 + \widehat{f}_a(0) + \int_0^\infty (p_a(t) - f_a(t))dt\right| \\ &\leq |F(0)| + \|p_a - f_a\|_{L^1} < 0 + \frac{\epsilon}{2} = \frac{\epsilon}{2}. \end{aligned}$$

Thus

$$\|F - P\|_{W^+} = \|f_a - p_a\|_{L^1} + \left|f_0 + \int_0^\infty p_a(t)dt\right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

This completes the proof. \square

Theorem 4.2.

(a) Let $\mathfrak{M}_0 := \{F \in W^+(\mathbb{C}_+) \mid F(0) = 0\}$ and

$$E_n := \frac{s}{s + \frac{1}{n}}, \quad n \in \mathbb{N}.$$

Then $(E_n)_{n \in \mathbb{N}}$ is a bounded approximate identity for \mathfrak{M}_0 .

(b) Let $\mathfrak{M}_\infty = \widehat{L_1(0, \infty)}$ and

$$U_n = \widehat{n\mathbf{1}_{[0, \frac{1}{n}]}}, \quad n \in \mathbb{N},$$

where $\mathbf{1}_{[0, \frac{1}{n}]}(t)$ is 1 if $t \in [0, \frac{1}{n}]$, and 0 otherwise. Then $(U_n)_{n \geq 1}$ is a bounded approximate identity for \mathfrak{M}_∞ .

Proof. (b) The existence of a bounded approximate identity for \mathfrak{M}_∞ follows [1, Theorem 6.5, p. 105]. The above example is easy to check.

(a) We note that

$$\|E_n\|_{W^+} = \left\| 1 + \widehat{\left(-\frac{1}{n}e^{-\frac{t}{n}}\right)} \right\|_{W^+} = |1| + \left\| -\frac{1}{n}e^{-\frac{t}{n}} \right\|_{L^1} = 1 + 1 = 2,$$

and so the sequence is bounded.

Given $F \in W^+(\mathbb{C}_+)$, and $\epsilon > 0$ arbitrarily small, in view of Lemma 4.1, we can find a $P \in \mathfrak{M}_0$ such that $P = \widehat{p_a} + p_0$, where $p_a \in L^1(0, \infty)$ has compact support, $p_0 \in \mathbb{C}$, and $\|F - P\|_{W^+} < \epsilon$. Then

$$\|E_n F - F\|_{W^+} \leq \|E_n P - P\|_{W^+} + \|E_n\|_{W^+} \|F - P\|_{W^+} + \|F - P\|_{W^+}.$$

So it is enough to prove that

$$\lim_{n \rightarrow \infty} \|E_n P - P\|_{W^+} = 0$$

for all $P \in \mathfrak{M}_0$ such that $P = \widehat{p_a} + p_0$, where $p_a \in L^1(0, \infty)$ has compact support, and $p_0 \in \mathbb{C}$. We do this below.

We have

$$E_n P - P = \frac{s + \frac{1}{n} - \frac{1}{n}}{s + \frac{1}{n}} P - P = -\frac{1}{n} \frac{1}{s + \frac{1}{n}} P = -\frac{1}{n} \left(\widehat{(e^{-t/n} * p_a)} + p_0 \widehat{e^{-t/n}} \right).$$

Let c be given by

$$c(t) := \int_0^t e^{-\frac{t-\tau}{n}} p_a(\tau) d\tau + p_0 e^{-\frac{t}{n}}.$$

Then $c \in L^1(0, \infty)$. Let $T > 0$ be such that $\text{supp}(p_a) \subset [0, T]$. We have

$$\|E_n P - P\|_{W^+} = \frac{1}{n} \|c\|_{L^1} = \frac{1}{n} \int_0^\infty |c(t)| dt = \underbrace{\frac{1}{n} \int_0^T |c(t)| dt}_{(I)} + \underbrace{\frac{1}{n} \int_T^\infty |c(t)| dt}_{(II)}.$$

We estimate (I) as follows:

$$\begin{aligned} (I) &= \frac{1}{n} \int_0^T |c(t)| dt = \frac{1}{n} \int_0^T \left| \int_0^t e^{-\frac{t-\tau}{n}} p_a(\tau) d\tau + p_0 e^{-\frac{t}{n}} \right| dt \\ &\leq \frac{1}{n} \int_0^T \left[\int_0^t e^{-\frac{t-\tau}{n}} |p_a(\tau)| d\tau + |p_0| e^{-\frac{t}{n}} \right] dt \\ &\leq \underbrace{\frac{1}{n} \int_0^T \left[\int_0^t 1 \cdot |p_a(\tau)| d\tau + |p_0| \cdot 1 \right] dt}_{(III)}. \end{aligned}$$

Since the integral (III) does not depend on n , we obtain that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \int_0^T |c(t)| dt = 0.$$

Furthermore,

$$\begin{aligned}
(II) &= \frac{1}{n} \int_T^\infty |c(t)| dt = \frac{1}{n} \int_T^\infty e^{-\frac{t}{n}} \left| \int_0^t e^{\frac{\tau}{n}} p_a(\tau) d\tau + p_0 \right| dt \\
&= \frac{1}{n} \int_T^\infty e^{-\frac{t}{n}} \left| \int_0^\infty e^{\frac{\tau}{n}} p_a(\tau) d\tau + p_0 \right| dt \quad (\text{since } \text{supp}(p_a) \subset [0, T]) \\
&= \frac{1}{n} \int_T^\infty e^{-\frac{t}{n}} \left| \widehat{p}_a \left(-\frac{1}{n} \right) + p_0 \right| dt
\end{aligned}$$

Since p_a has compact support in $[0, T]$, \widehat{p}_a is an entire function by the Paley-Wiener theorem; see for instance [11, Theorem 7.2.3, p. 122]. Consequently,

$$\begin{aligned}
(II) &= \frac{1}{n} \int_T^\infty e^{-\frac{t}{n}} \left| \widehat{p}_a \left(-\frac{1}{n} \right) + p_0 \right| dt \\
&= \frac{1}{n} \int_T^\infty e^{-\frac{t}{n}} dt \cdot \left| \widehat{p}_a \left(-\frac{1}{n} \right) + p_0 \right| \\
&= e^{-\frac{T}{n}} \left| \widehat{p}_a \left(-\frac{1}{n} \right) + p_0 \right| \xrightarrow{n \rightarrow \infty} 1 \cdot |\widehat{p}_a(0) + p_0| = |P(0)| = 0.
\end{aligned}$$

This completes the proof of the case (a). \square

The case of $\mathfrak{M}_{i\beta}$ works in a similar manner.

Theorem 4.3. *Let $F \in W^+(\mathbb{C}_+)$, and let $s_0 \in \mathbb{C}$ be such that $\text{Re}(s_0) > 0$ and $F(s_0) = 0$. Then $\frac{F}{s-s_0} \in W^+(\mathbb{C}_+)$.*

Proof. Let $F = \widehat{f}_a + f_0$, where $f_a \in L^1(0, \infty)$ and $f_0 \in \mathbb{C}$. Since $F(s_0) = 0$, we have

$$(1) \quad F(s_0) = \int_0^\infty e^{-s_0\tau} f_a(\tau) d\tau + f_0 = 0.$$

Let c be defined by

$$(2) \quad c(t) = \begin{cases} f_0 e^{s_0 t} + \int_0^t e^{s_0(t-\tau)} f_a(\tau) d\tau & \text{for } t > 0, \\ 0 & \text{for } t < 0. \end{cases}$$

We have

$$\begin{aligned}
\int_0^\infty |c(t)|dt &= \int_0^\infty e^{\operatorname{Re}(s_0)t} \left| f_0 + \int_0^t e^{-s_0\tau} f_a(\tau) d\tau \right| dt \\
&= \int_0^\infty e^{\operatorname{Re}(s_0)t} \left| - \int_t^\infty e^{-s_0\tau} f_a(\tau) d\tau \right| dt \quad (\text{using (1)}) \\
&\leq \int_0^\infty e^{\operatorname{Re}(s_0)t} \int_t^\infty e^{-\operatorname{Re}(s_0)\tau} |f_a(\tau)| d\tau dt \\
&= \int_0^\infty \int_t^\infty e^{\operatorname{Re}(s_0)t} e^{-\operatorname{Re}(s_0)\tau} |f_a(\tau)| d\tau dt \\
&= \int_0^\infty \int_0^\tau e^{\operatorname{Re}(s_0)t} e^{-\operatorname{Re}(s_0)\tau} |f_a(\tau)| dt d\tau \\
&= \int_0^\infty e^{-\operatorname{Re}(s_0)\tau} |f_a(\tau)| \int_0^\tau e^{\operatorname{Re}(s_0)t} dt d\tau \\
&= \int_0^\infty e^{-\operatorname{Re}(s_0)\tau} |f_a(\tau)| \frac{e^{\operatorname{Re}(s_0)\tau} - 1}{\operatorname{Re}(s_0)} d\tau \\
&\leq \frac{1}{\operatorname{Re}(s_0)} \int_0^\infty |f_a(\tau)| d\tau < \infty.
\end{aligned}$$

So $c \in L^1(0, \infty)$.

Let $\beta \in \mathbb{C}$ be such that $\operatorname{Re}(\beta) > \operatorname{Re}(s_0)$. Then we have from (2) that

$$(3) \quad e^{-\beta t} c(t) = \begin{cases} f_0 e^{(s_0-\beta)t} + \left([e^{(s_0-\beta)x} u] * [e^{-\beta x} f_a] \right)(t) & \text{for } t > 0 \\ 0 & \text{for } t < 0 \end{cases}$$

where u denotes the step function, given by $u(t) = 1$ for $t > 0$ and $u(t) = 0$ otherwise.

Recall the fact that if $g_a \in L^1(0, \infty)$, then for a complex number α such that $\operatorname{Re}(\alpha) > 0$, $(\widehat{e^{-\alpha t} g_a})(s) = \widehat{g_a}(s + \alpha)$ (for $s \in \mathbb{C}_+$). Using this, we obtain

$$(\widehat{e^{-\beta t} c})(s) = \widehat{c}(s + \beta) \quad \text{and} \quad (\widehat{e^{-\beta t} f_a})(s) = \widehat{f_a}(s + \beta) \quad (s \in \mathbb{C}_+).$$

Since the Laplace transform of a convolution is the product of the Laplace transforms (see for instance [4, Proposition 14.1]), we have

$$\left(\widehat{[e^{(s_0-\beta)x} u] * [e^{-\beta x} f_a]} \right)(s) = \frac{1}{s + \beta - s_0} \cdot \widehat{f_a}(s + \beta) \quad (s \in \mathbb{C}_+).$$

Using these facts, we see by taking Laplace transform on both sides of (3) that

$$\widehat{c}(s + \beta) = \frac{f_0}{s + \beta - s_0} + \frac{1}{s + \beta - s_0} \cdot \widehat{f_a}(s + \beta) = \frac{F(s + \beta)}{s + \beta - s_0} \quad (s \in \mathbb{C}_+).$$

So for all s such that $\operatorname{Re}(s) > \operatorname{Re}(s_0)$, we have

$$\widehat{c}(s) = \frac{F(s)}{s - s_0}.$$

By the identity principle, the above holds in \mathbb{C}_+ . So $\frac{F}{s-s_0} = \widehat{c} \in W^+(\mathbb{C}_+)$. \square

In our proof of Theorem 1.5 will need the corona theorem for $W^+(\mathbb{C}_+)$:

Proposition 4.4. *If $F_1, \dots, F_n \in W^+(\mathbb{C}_+)$ are such that there exists a $\delta > 0$ such that*

$$(4) \quad \text{for all } s \in \mathbb{C}_+, \quad |F_1(s)| + \dots + |F_n(s)| > \delta > 0,$$

then there exist $G_1, \dots, G_n \in W^+(\mathbb{C}_+)$ such that

$$(5) \quad \text{for all } s \in \mathbb{C}_+, \quad G_1(s)F_1(s) + \dots + G_n(s)F_n(s) = 1.$$

Proof. That (5) implies (4) is easy to see. The reverse implication follows from the classical result (see [3, p.112]) that the maximal ideals of $W^+(\mathbb{C}_+)$ are given by \mathfrak{M}_∞ and \mathfrak{M}_{s_0} , where $s_0 \in \mathbb{C}_+$. Indeed, suppose that $F_1, \dots, F_n \in W^+(\mathbb{C}_+)$ satisfy (4), but that the ideal $(F_1, \dots, F_n) \neq (1)$. Then there exists a maximal ideal \mathfrak{M} that contains (F_1, \dots, F_n) . We now consider the two possible cases:

- (i) If $\mathfrak{M} = \mathfrak{M}_{s_0}$ for some $s_0 \in \mathbb{C}_+$, then $(F_1, \dots, F_n) \subset \mathfrak{M}_{s_0}$ yields that $F_1(s_0) = \dots = F_n(s_0) = 0$, which contradicts (4).
- (ii) Now suppose that $\mathfrak{M} = \mathfrak{M}_\infty$. Let $F_k = \widehat{f_{k,a}} + f_{k,0}$, where $f_{k,a} \in L^1(0, \infty)$ and $f_{k,0} \in \mathbb{C}$, $k = 1, \dots, n$. Since $(F_1, \dots, F_n) \subset \mathfrak{M}_\infty$, we have $f_{1,0} = \dots = f_{n,0} = 0$. Hence $F_k = \widehat{f_{k,a}}$, $k = 1, \dots, n$. Passing to the limit $s \rightarrow \infty$ in (4), we obtain the contradiction that $0 \geq \delta$.

Consequently $(F_1, \dots, F_n) = (1)$, and so (5) holds for some $G_1, \dots, G_n \in W^+(\mathbb{C}_+)$. \square

Lemma 4.5. *Suppose that $F_1, \dots, F_n \in W^+(\mathbb{C}_+)$ and that D is a greatest common divisor of F_1, \dots, F_n . If $s_0 \in \mathbb{C}_+$ is a common zero of F_1, \dots, F_n , then $D(s_0) = 0$ as well.*

Proof. If $\operatorname{Re}(s_0) > 0$, then let m be the least integer among the multiplicities of s_0 as a zero respectively of F_1, \dots, F_n . By Theorem 4.3, $(s - s_0)^m$ is a divisor of F_1, \dots, F_n . But since D is the greatest common divisor of F_1, \dots, F_n , it follows that $(s - s_0)^m$ divides D .

If on the other hand $\operatorname{Re}(s_0) = 0$, then $F_1, \dots, F_n \in \mathfrak{M}_{s_0}$, where

$$\mathfrak{M}_{s_0} := \{F \in W^+(\mathbb{C}_+) \mid F(s_0) = 0\}.$$

By Theorem 4.2, \mathfrak{M}_{s_0} has a bounded approximate identity. Applying Proposition 2.2, with $(a_n)_{n \geq 1} := (F_1, \dots, F_n, 0, 0, 0, \dots)$, we get the existence of an element $C \in \mathfrak{M}_{s_0}$, and $G_1, \dots, G_n \in \mathfrak{M}_{s_0}$ such that $F_k = CG_k$, $k = 1, \dots, n$. So we have a common divisor C of F_1, \dots, F_n . Since D is a greatest common divisor, C divides D and so $D(s_0) = 0$, too. \square

Lemma 4.6. *Suppose that $F_1, \dots, F_n \in \mathfrak{M}_\infty$ and that $D \in W^+(\mathbb{C}_+)$ is a greatest common divisor of F_1, \dots, F_n . Then $D \in \mathfrak{M}_\infty$ as well.*

Proof. Applying Theorem 4.2 and Proposition 2.2 to the sequence $(a_n) = (F_1, F_2, \dots, F_n, 0, 0, 0 \dots)$ we get a common divisor $C \in \mathfrak{M}_\infty$ of the F_j 's, $j = 1, \dots, n$. Hence C divides D ; that is $D = QC$ for some $Q \in W^+(\mathbb{C}_+)$. Therefore $D \in \mathfrak{M}_\infty$. \square

Proof of Theorem 1.5. Let $F_1, \dots, F_n \in W^+(\mathbb{C}_+)$ have a greatest common divisor D . By the algebraic result in Lemma 2.3, it follows that 1 is a greatest common divisor of $\frac{F_1}{D}, \dots, \frac{F_n}{D}$.

Lemma 4.5 gives

$$(6) \quad \left| \frac{F_1}{D} \right| + \dots + \left| \frac{F_n}{D} \right| > 0 \text{ in } \mathbb{C}_+.$$

Since $F_k/D \in W^+(\mathbb{C}_+)$ for each k , $F_k/D = \widehat{h_k} + \alpha_k$, where $h_k \in L^1(0, \infty)$ and $\alpha_k \in \mathbb{C}$. Since $h_k \in L^1(0, \infty)$, we have

$$(7) \quad \lim_{\substack{s \rightarrow \infty \\ s \in \mathbb{C}_+}} \widehat{h_k}(s) = 0.$$

We consider the two possible cases:

- (i) All the α_k 's are zero. Then by Lemma 4.6, it follows that 1 is the Laplace transform of an element in $L^1(0, \infty)$, which is a contradiction.
- (ii) At least one of the α_k 's is not zero. Then $|\alpha_1| + \dots + |\alpha_n| > 0$. So for $s \in \mathbb{C}_+$ such that $|s| > R$ with a large enough R , (7) gives the existence of a $\delta > 0$ such that

$$\left| \frac{F_1}{D} \right| + \dots + \left| \frac{F_n}{D} \right| > \delta > 0,$$

while on the compact set K consisting of $s \in \mathbb{C}_+$ with $|s| \leq R$, $\left| \frac{F_1}{D} \right| + \dots + \left| \frac{F_n}{D} \right|$ is at least as large as its minimum value on K , which is positive by (6). So by Proposition 4.4, it follows that there exist $G_1, \dots, G_n \in W^+(\mathbb{C}_+)$ such that

$$G_1 \frac{F_1}{D} + \dots + G_n \frac{F_n}{D} = 1,$$

and so $G_1 F_1 + \dots + G_n F_n = D$.

This completes the proof of the theorem. \square

REFERENCES

- [1] L.W. Baggett. *Functional Analysis: A Primer*. Marcel Dekker, New York, 1992.
- [2] V.M. Faïvševskii. The structure of the ideals of certain algebras of analytic functions. (Russian) *Dokl. Akad. Nauk SSSR*, 211:537-539, 1973. English translation in *Soviet Math. Dokl.*, 14:1067-1070, 1973.
- [3] I. Gelfand, D. Raikov, G. Shilov. *Commutative Normed Rings*. Chelsea Publ. Comp. New York, 1964.
- [4] P.B. Guest. *Laplace Transforms and an Introduction to Distributions*. Ellis Norwood, 1991.

- [5] K. Koua. Un exemple d'algèbre de Banach commutative radicale a unité approchée bornée sans multiplicateur non trivial. *Mathematica Scandinavica*, no. 1, 56:70-82, 1985.
- [6] H. Logemann. Finitely generated ideals in certain algebras of transfer functions for infinite-dimensional systems. *International Journal of Control*, no. 1, 45:247-250, 1987.
- [7] R. Mortini. The Chang-Marshall algebras. *Mitteilungen aus dem Mathematischen Seminar Giessen*, no. 185:1-76, 1988.
- [8] R. Mortini and M. von Renteln. Ideals in the Wiener algebra W^+ . *Journal of the Australian Mathematical Society (Series A)*, 46: 220-228, 1990.
- [9] A. Quadrat. An Introduction to Control Theory. Course Notes, *Mathematics, Algorithms, Proofs*, Castro Urdiales (Spain), 09-13 January, 2006. Available at www-sop.inria.fr/cafe/Alban.Quadrat/Stage/MAp1.pdf.
- [10] A. Quadrat. The fractional representation approach to synthesis problems: an algebraic analysis viewpoint. Part I: (Weakly) doubly coprime factorizations. *SIAM Journal on Control and Optimization*, 42:266-299, 2004.
- [11] R.S. Strichartz. *A Guide to Distribution Theory and Fourier Transforms*. World Scientific, Singapore, 2003.
- [12] M. von Renteln. Divisibility structure and finitely generated ideals in the disc algebra. *Monatshefte für Mathematik*, 82:51-56, 1976.
- [13] M. von Renteln. A simple constructive proof of an analogue of the corona theorem. *Proceedings of the American Mathematical Society*, 83:299-303, no. 2, 1981.
- [14] N. Th. Varopoulos. Sur les formes positives d'une algèbre de Banach. (French) *Comptes Rendus Acad. Sci. Paris* 258:2465-2467, Groupe 1, 1964.

MORTINI: UNIVERSITÉ PAUL VERLAINE - METZ, LMAM ET DÉPARTEMENT DE MATHÉMATIQUES, ILE DU SAULCY, F-57045 METZ FRANCE.

E-mail address: `mortini@univ-metz.fr`

SASANE: MATHEMATICS DEPARTMENT, LONDON SCHOOL OF ECONOMICS, HOUGHTON STREET, LONDON WC2A 2AE, UNITED KINGDOM.

E-mail address: `A.J.Sasane@lse.ac.uk`